

COMPARISONS BETWEEN PERIODIC, ANALYTIC AND LOCAL CYCLIC COHOMOLOGY

RALF MEYER

ABSTRACT. We compute periodic, analytic and local cyclic cohomology for convolution algebras of compact Lie groups in order to exhibit differences between these theories. A surprising result is that the periodic and analytic cyclic cohomology of the smooth convolution algebras differ, although these algebras have finite homological dimension.

1. INTRODUCTION

Cyclic cohomology is a non-commutative generalization of de Rham homology. Besides cyclic cohomology itself, there are several variants like periodic cyclic cohomology, entire cyclic cohomology and local cyclic cohomology (see [4, 5, 12]). As in [9], we use the more appropriate name “analytic cyclic cohomology” instead of “entire cyclic cohomology” in this note. We denote these theories by HP^* (periodic), HA^* (analytic) and HL^* (local), respectively. We compute them—together with the dual homology theories—for some convolution algebras of compact Lie groups. We consider these apparently trivial examples because they clearly exhibit differences between the three theories.

Throughout this note, K is a compact Lie group. Since we aim for counterexamples, it would be sufficient to consider only the circle group S^1 . We allow general compact Lie groups because this creates no additional difficulties. Let $C^*(K)$ be the group C^* -algebra of K , and let $C^\infty(K)$ and $\mathcal{H}(K)$ be the dense subalgebras of smooth functions and of coefficients of finite dimensional representations of K , respectively. For $K = S^1$, we get

$$C^*(S^1) \cong C_0(\mathbb{Z}), \quad C^\infty(S^1) \cong \mathcal{S}(\mathbb{Z}), \quad \mathcal{H}(K) \cong \bigoplus_{n \in \mathbb{Z}} \mathbb{C},$$

where the product on the right hand sides is the pointwise one and $\mathcal{S}(\mathbb{Z})$ is the space of rapidly decreasing sequences.

It follows from the general properties of the local theory that the natural maps

$$HL_*(\mathcal{H}(K)) \rightarrow HL_*(C^\infty(K)) \rightarrow HL_*(C^*(K)) \leftarrow K_*(C^*(K)) \otimes_{\mathbb{Z}} \mathbb{C}$$

are all isomorphisms. The last map is the Chern-Connes character. A similar assertion holds in cohomology. Thus the local theory provides a perfect description of the de Rham (co)homology of the space of representations $\hat{K}$.

In order to show that the Chern-Connes character above is an isomorphism, we prove a Universal Coefficient Theorem for the bivariant local cyclic cohomology of C^* -algebras. If A and B are C^* -algebras that satisfy the UCT in bivariant Kasparov theory, then there is a natural isomorphism

$$HL(A, B) \cong \text{Hom}(K_*(A) \otimes_{\mathbb{Z}} \mathbb{C}, K_*(B) \otimes_{\mathbb{Z}} \mathbb{C})$$

of graded complex vector spaces.

2000 *Mathematics Subject Classification.* 19D55, 18G60.

This research was supported by the EU-Network *Quantum Spaces and Noncommutative Geometry* (Contract HPRN-CT-2002-00280) and the *Deutsche Forschungsgemeinschaft* (SFB 478).

For the very small subalgebra $\mathcal{H}(K)$, all three cyclic theories still agree, both in cohomology and homology. This follows easily from their behavior for direct sums, which we investigate in Section 3.

For $C^\infty(K)$ the computation of Hochschild, cyclic and periodic cyclic cohomology is still straightforward because $C^\infty(K)$ is biprojective. This implies also that its homological dimension is at most 2. It is remarkable that the natural maps

$$\mathrm{HP}^*(\mathcal{H}(K)) \leftarrow \mathrm{HP}^*(C^\infty(K)), \quad \mathrm{HP}_*(\mathcal{H}(K)) \rightarrow \mathrm{HP}_*(C^\infty(K))$$

fail to be isomorphisms. Thus the periodic theory is surprisingly sensitive to the choice of a “smooth” subalgebra. If we want to describe the de Rham homology of $\hat{K}$ by periodic cyclic cohomology, the subalgebra $C^\infty(K)$ is not yet small enough. We have to go down further to $\mathcal{H}(K)$.

General properties of the theories show that analytic and local cyclic homology coincide for our three examples. In cohomology such a result is not available. At the moment, direct computations of analytic cyclic cohomology seem to be impossible. The only available general method to compute it is to use its homological properties, in particular, six term exact sequences. However, since the periodic and analytic theory have the same homological properties, this method can only work for algebras for which the two theories agree. Nevertheless, we can use the Chern-Connes character in K-homology to construct non-trivial analytic cyclic cocycles. We can even show that the natural map from the analytic to the local cyclic cohomology is surjective both for $C^\infty(K)$ and $C^*(K)$.

As a result, analytic and periodic cyclic (co)homology must differ for $C^\infty(K)$, although this algebra has finite homological dimension. Hence Khalkhali’s result that periodic and analytic cyclic cohomology agree for a Banach algebra of finite homological dimension fails for these very elementary nuclear Fréchet algebras.

2. PREPARATIONS

We do not include the definitions of the cyclic theories we are considering because that would take too much space. We refer the reader to [4, 5, 8, 9, 12]. All cyclic cohomology theories are defined for algebras with additional structure like a topology or bornology. For the analytic and local theory, it is essential to use either bornological algebras or inductive systems. Since this setup may be unfamiliar to many readers we briefly explain what bornological vector spaces are. See [9] for further details and references.

A complete bornological vector space is a vector space V together with a *bornology*, which is a collection $\mathfrak{S}(V)$ of subsets that are called *bounded*. The prototypical example of a bornology is the family of “bounded” subsets of a (quasi)complete locally convex topological vector space, which we call the *von Neumann bornology*. We equip the smooth convolution algebra $C^\infty(K)$ with this bornology. Thus a subset $S \subseteq C^\infty(K)$ is bounded if and only if the functions $D(f): K \rightarrow \mathbb{C}$ for $f \in S$ are uniformly bounded for any differential operator D on K .

Another example is the *fine bornology*, which is defined on any vector space. It consists of the bounded subsets of finite dimensional subspaces. We equip the algebra $\mathcal{H}(K)$ with this bornology. Notice that the topology of $\mathcal{H}(K)$ is much more complicated to describe than its bornology.

For local cyclic cohomology it is important to choose a smaller bornology than the von Neumann bornology. If A is a Banach algebra or a C^* -algebra, we *always* equip it with the *precompact bornology*, which consists of precompact subsets of A . In a complete space a subset is precompact if and only if it is relatively compact if and only if it is contained in a compact subset. In particular, this is the bornology we choose on $C^*(K)$. For the Montel space $C^\infty(K)$, the von Neumann bornology

and the precompact bornology coincide. Otherwise, we should not have chosen the von Neumann bornology on $C^\infty(K)$.

We have to use the precompact bornology because approximation results in Banach spaces usually hold uniformly on precompact subsets, but not on von Neumann bounded subsets. For instance, the identity map on a Banach space with Grothendieck's approximation property can be approximated uniformly on *precompact* sets by finite rank maps. Similarly, an asymptotic morphism is approximately multiplicative uniformly on *precompact* subsets, but usually not on von Neumann bounded subsets.

Since we use the precompact instead of the von Neumann bornology in the definition of analytic cyclic cohomology for Banach algebras, we get more analytic cocycles than in the usual definition of entire cyclic cohomology in [5]. Our computations also show that this change of bornology has a drastic effect on the resulting theory.

The right morphisms between bornological vector spaces are the bounded maps. Moreover, we have an obvious notion of a bounded (bi)linear map. A bornological algebra is a bornological vector space with a bounded associative multiplication. It is straightforward to verify that $\mathcal{H}(K)$, $C^\infty(K)$ and $C^*(K)$ with the bornologies defined above are complete bornological algebras.

The completed bornological tensor product $\hat{\otimes}$ is defined by a universal property for bounded bilinear maps. We use this completed tensor product to construct the complexes computing the various homology theories. For our examples we have isomorphisms of bornological vector spaces

$$\mathcal{H}(K) \hat{\otimes} \mathcal{H}(K) \cong \mathcal{H}(K \times K), \quad C^\infty(K) \hat{\otimes} C^\infty(K) \cong C^\infty(K \times K).$$

The space $C^*(K) \hat{\otimes} C^*(K)$ is more complicated.

The periodic cyclic theory is usually defined for topological algebras. It is shown in [9] that for Fréchet algebras with the precompact bornology, our bornological approach yields the same Hochschild, cyclic and periodic cyclic (co)homology groups as the topological approach. We even get the same complexes. This equivalence between topology and bornology is based on Grothendieck's results about compact subsets of projective tensor products.

The bornological approach is much more flexible than the topological approach. There seems to be no convolution algebra on a compact Lie group which is a Fréchet algebra and yields the correct periodic cyclic cohomology. We have to use $\mathcal{H}(K)$, which is best viewed as a bornological algebra (or as an algebra without additional structure).

A separately continuous bilinear map defined on (quasi)complete topological vector spaces is automatically bounded. Hence any quasi-complete locally convex algebra with separately continuous multiplication becomes a bornological algebra when equipped with the von Neumann or precompact bornology. Therefore, the bornological tensor product is similar to the completed injective tensor product for topological vector spaces, which is universal for separately continuous bilinear maps. However, it enjoys much better properties because boundedness is a more tractable condition than separate continuity. For instance, the bornological tensor product is associative, the *completed* injective tensor product is only associative under additional assumptions.

Studying bounded subsets of a bornological vector space means that we approximate a bornological vector space from within by Banach spaces. In fact, the basic structure theorem about complete bornological vector spaces asserts that each such space can be written in a canonical way as an inductive limit of Banach spaces. Furthermore, this construction identifies the category of complete bornological vector spaces with a full subcategory of the category of inductive systems of Banach

spaces. The inductive systems that come from bornological vector spaces have the additional property that the structure maps are all injective.

For the homological algebra behind the local cyclic theory, it is useful to allow arbitrary inductive systems. Therefore, Puschnigg defines local cyclic cohomology on the category of inductive systems of Banach algebras. He also allows inductive systems of *admissible Fréchet algebras*. A Fréchet algebra is called admissible if and only if there is an open neighborhood of zero such that for any compact subset $S \subset U$, its *multiplicative closure* $\bigcup_{n=1}^{\infty} S^n$ is again precompact. For instance, Banach algebras and smooth subalgebras of Banach algebras (in the sense of Puschnigg [12]) are admissible Fréchet algebras. However, since any admissible Fréchet algebra can be written as an inductive limit of Banach algebras, this seemingly greater generality only makes the notation more complicated.

We remark that a complete bornological algebra need not be an inductive limit of Banach *algebras* because there may be no multiplicatively closed bounded subsets. For instance, the group ring of an infinite discrete group is never an inductive limit of Banach algebras. It is possible, but technically difficult, to extend Puschnigg's theory to the larger category of algebra objects in the category of inductive systems of Banach spaces. It is easier to transport local cohomology to the bornological category. This approach yields equivalent results in all relevant applications. We will show this in a forthcoming article.

Fortunately, the bornological algebras that we have to consider are all inductive limits of Banach algebras. The Peter-Weyl theorem asserts that $\mathcal{H}(K)$ is a countable direct sum of matrix algebras, the sum being indexed by the irreducible representations of K . The algebras $C^\infty(K)$ and $C^*(K)$ are even admissible Fréchet algebras because the multiplicative closure of any compact subset of the open unit ball of $C^*(K)$ is again precompact.

We will stay away from the details of the machinery of local cyclic cohomology, where inductive systems play a rôle. As a result, we may just as well work in the more intuitive setup of bornological algebras.

3. BEHAVIOR FOR DIRECT SUMS

Both for the computation of the theories for $\mathcal{H}(K)$ and for the proof of the Universal Coefficient Theorem for the local cyclic theory, we need the compatibility of the various cyclic theories with (countable) direct sums. The direct sum in the category of bornological algebras is just the algebraic direct sum equipped with the coarsest bornology making the inclusions of the summands bounded. Hence a subset of $\bigoplus_{j \in J} A_j$ is bounded if and only if it is contained in and bounded in a finite sum $\bigoplus_{j \in F} A_j$ for some finite subset $F \subseteq J$.

Proposition 3.1. *Let $(A_j)_{j \in \mathbb{N}}$ be a sequence of complete bornological algebras and let $A := \bigoplus A_j$ be the bornological direct sum. Let B be another bornological algebra. Then we have:*

$$\begin{aligned} \mathrm{HH}_*(A) &\cong \bigoplus \mathrm{HH}_*(A_j), & \mathrm{HH}^*(A) &\cong \prod \mathrm{HH}^*(A_j), \\ \mathrm{HC}_*(A) &\cong \bigoplus \mathrm{HC}_*(A_j), & \mathrm{HC}^*(A) &\cong \prod \mathrm{HC}^*(A_j), \\ \mathrm{HA}_*(A) &\cong \bigoplus \mathrm{HA}_*(A_j), & \mathrm{HA}^*(A, B) &\cong \prod \mathrm{HA}^*(A_j, B), \\ \mathrm{HL}_*(A) &\cong \bigoplus \mathrm{HL}_*(A_j), & \mathrm{HL}^*(A, B) &\cong \prod \mathrm{HL}^*(A_j, B), \end{aligned}$$

For the local theory, we have to assume all A_j and B to be inductive limits of Banach algebras in order for the theory to be defined.

Proof. All assertions are easy to obtain for *finite* sums. For the analytic and local cyclic theory, this additivity follows from excision. Although Hochschild and cyclic

(co)homology do not satisfy excision in complete generality, they do satisfy it for direct sum extensions (see [8]). We will reduce countable sums to finite sums.

The Hochschild, cyclic and analytic cyclic (co)homology of A are all defined as the (co)homology of certain complexes $C(A)$ associated to the algebra A in a natural way. For the local theory, we have to take the “local cohomology” of the same complex as for the analytic theory, viewed as an inductive system of complexes. However, since local cohomology enjoys even better properties than ordinary cohomology, this is only a notational change. For the bivariant theories, we have to consider morphisms $C(A) \rightarrow C(B)$ in an appropriate (local) homotopy category. Again this only creates notational problems. What we really have to show is that $C(A)$ is chain homotopic to the direct sum of the complexes $C(A_j)$ for all theories we consider. We will use that this is true for finite sums. The proof of additivity for finite sums actually yields this stronger statement in all cases.

We will not define the complexes $C(A)$ here because we only use simple formal properties. Besides naturality and the result for finite sums, we need the following compatibility with inductive limits: if (A'_n) is a sequence of bornological algebras with injective structure maps $A'_n \rightarrow A'_{n+1}$, then $C(\varinjlim A'_n) = \varinjlim C(A'_n)$.

We apply this to $A'_n := \bigoplus_{j=1}^n A_j$. Since A is the bornological direct limit of (A'_n) , we conclude that $C(A)$ is the bornological direct limit of the complexes $C(A'_n)$ for $n \rightarrow \infty$. Since A'_{n-1} is a retract of A'_n , the complex $C(A'_{n-1})$ is a direct summand of $C(A'_n)$. Write $C(A'_n) = C(A'_{n-1}) \oplus X_n$. Then we have $C(A'_n) = X_1 \oplus \cdots \oplus X_n$. As a result, $C(A) \cong \bigoplus X_n$. Additivity for finite sums implies that the complexes $C(A_n)$ and X_n are chain homotopic. Thus $C(A)$ is chain homotopic to the direct sum $\bigoplus C(A_j)$. The assertions of the proposition follow. $\square$

We do not have a general assertion about direct sums in the second variable of the analytic and local cyclic theories. This should be expected because direct sums and products do not commute.

It is remarkable that the periodic theory may behave quite badly with respect to direct sums. The periodic cyclic cohomology is isomorphic to the inductive limit of the cyclic cohomology groups with respect to the periodicity operator S . Since the cyclic cohomology of a direct sum is a direct product, the periodic cyclic cohomology of a direct sum is an inductive limit of direct products. However, direct products and inductive limits do not commute!

4. THE UNIVERSAL COEFFICIENT THEOREM

The local cyclic theory has very good functorial properties on the category of C^* -algebras. This is the source of the Universal Coefficient Theorem (UCT) below. It suffices to state it for the bivariant theory because

$$\mathrm{HL}_*(A) := \mathrm{HL}_*(\mathbb{C}, A), \quad \mathrm{HL}^*(A) := \mathrm{HL}_*(A, \mathbb{C}).$$

Recall that C^* -algebras are always equipped with the precompact bornology.

Theorem 4.1 (Universal Coefficient Theorem). *Let A and B be separable C^* -algebras that satisfy the Universal Coefficient Theorem in Kasparov’s KK-theory. Thus A and B are KK-equivalent to commutative C^* -algebras.*

Then there are natural isomorphisms

$$\mathrm{HL}(A, B) \cong \mathrm{Hom}(\mathrm{HL}_*(A), \mathrm{HL}_*(B)) \cong \mathrm{Hom}(K_*(A) \otimes_{\mathbb{Z}} \mathbb{C}, K_*(B) \otimes_{\mathbb{Z}} \mathbb{C}).$$

Proof. First we prove that the Chern-Connes character is an isomorphism

$$(1) \quad K_*(B) \otimes_{\mathbb{Z}} \mathbb{C} \cong \mathrm{KK}_*(\mathbb{C}, B) \otimes_{\mathbb{Z}} \mathbb{C} \cong \mathrm{HL}_*(\mathbb{C}, B) = \mathrm{HL}_*(B)$$

provided that B satisfies the UCT in KK-theory. This already proves that the second map in the theorem is an isomorphism. There is a bivariant Chern-Connes

character $\mathrm{KK} \rightarrow \mathrm{HL}$ because HL is a stable exact homotopy functor in both variables and KK -theory is universal among such theories. Invariance for continuous homotopies and C^* -stability for HL are proved in [12]. Excision is proved in [11] (see also [9, 10]).

Since a separable C^* -algebra satisfies the UCT if and only if it is KK -equivalent to a commutative C^* -algebra, it suffices to prove the assertion for B in the *bootstrap category* (see [2]). This is the smallest class of separable *nuclear* C^* -algebras that

- contains $\mathbb{C}$;
- is closed under KK -equivalence;
- is closed under countable direct sums;
- is closed under extensions in the following sense: if two C^* -algebras in an extension belong to the class, so does the third.

In [2], the third condition is replaced by the seemingly stronger requirement of being closed under countable inductive limits. However, a mapping telescope argument shows that inductive limits can be reduced to direct sums and pull backs and pull backs can be reduced to extensions (see [13]).

Hence it suffices to show that the class of separable nuclear C^* -algebras for which (1) holds has these properties. Since $\mathrm{HL}_0(\mathbb{C}, \mathbb{C}) = \mathbb{C}$ and $\mathrm{HL}_1(\mathbb{C}, \mathbb{C}) = 0$, it certainly contains $\mathbb{C}$. It is also closed under KK -equivalence by the existence of the bivariant Chern-Connes character. It is closed under countable direct sums because

$$\mathrm{HL}(\mathbb{C}, \bigoplus^{C^*} B_j) \cong \bigoplus \mathrm{HL}(\mathbb{C}, B_j)$$

for any sequence of C^* -algebras (B_j) . Here $\bigoplus^{C^*} B_j$ denotes the direct sum in the category of C^* -algebras. This direct sum is HL -equivalent to the bornological direct sum by Theorem 3.15 of [12]. Since the finite sums are retracts of the infinite sum, the technical assumption for that theorem is satisfied. Hence the claim follows from Proposition 3.1.

An extension of separable nuclear C^* -algebras automatically has a bounded, even completely positive, linear section. Hence it gives rise to six term exact sequences both in $K_* \otimes_{\mathbb{Z}} \mathbb{C}$ and HL_* . A diagram chase shows that our class is closed under extensions. As a result, (1) holds for all B in the bootstrap category.

Next we consider the class of separable nuclear C^* -algebras A for which the product in local cyclic cohomology gives rise to an isomorphism

$$\mathrm{HL}(A, B) \rightarrow \mathrm{Hom}(\mathrm{HL}_*(A), \mathrm{HL}_*(B))$$

for all bornological algebras B . We claim that this class contains the bootstrap category. The proof of this claim will complete the proof of the theorem.

The computation of $\mathrm{HL}_*(\mathbb{C})$ shows that $\mathbb{C}$ belongs to this class. Again the class is closed under KK -equivalence because of the existence of the bivariant Chern-Connes character. The same arguments as above treat direct sums and extensions. Hence the class contains the bootstrap category. $\square$

We remark that if the UCT is true, then the Chern-Connes character

$$\mathrm{KK}(A, B) \otimes_{\mathbb{Z}} \mathbb{C} \rightarrow \mathrm{HL}(A, B)$$

is always injective and has dense range in an appropriate sense. Nevertheless, it fails to be an isomorphism for $A = C^*(K)$ and $B = \mathbb{C}$.

5. COMPUTATION OF LOCAL CYCLIC COHOMOLOGY

Since $C^*(K)$ is a C^* -direct sum of matrix algebras, it belongs to the bootstrap category. Hence the UCT of the previous section applies. The K -theory of $C^*(K)$

vanishes in odd degrees and is isomorphic to the representation ring $R(K)$ of K in even degrees. Hence

$$\begin{aligned} \mathrm{HL}_0(C^*(K)) &\cong R(K) \otimes_{\mathbb{Z}} \mathbb{C}, & \mathrm{HL}_1(C^*(K)) &\cong \{0\}, \\ \mathrm{HL}^0(C^*(K)) &\cong \mathrm{Hom}_{\mathbb{Z}}(R(K), \mathbb{C}), & \mathrm{HL}^1(C^*(K)) &\cong \{0\}. \end{aligned}$$

Since $R(K)$ has a countable basis, we can identify $\mathrm{HL}_0(C^*(K))$ and $\mathrm{HL}^0(C^*(K))$ with the direct sum and the direct product of countably many copies of $\mathbb{C}$.

Proposition 5.1. *The inclusions $\mathcal{H}(K) \subseteq C^\infty(K) \subseteq C^*(K)$ are HL-equivalences and hence induce isomorphisms in local cyclic (co)homology.*

Proof. We claim that Theorem 3.2 in [12] applies to our situation. It yields that the inclusions $\mathcal{H}(K) \rightarrow C^*(K)$ and $\mathcal{H}(K) \rightarrow C^\infty(K)$ are isomorphisms in his “stable strongly almost multiplicative homotopy category.” Hence they are HL-equivalences and the proposition follows.

To apply Puschnigg’s result, observe that $\mathcal{H}(K)$ is a direct sum of finite dimensional matrix algebras and hence the increasing union of a sequence (V_n) of finite dimensional subalgebras. We have already remarked above that $C^*(K)$ and $C^\infty(K)$ are separable admissible Fréchet algebras. We also need to know that they possess Grothendieck’s approximation property. This is clear because $C^*(K)$ is nuclear as a C^* -algebra and $C^\infty(K)$ is nuclear as a locally convex vector space. $\square$

Thus the computation of the local theory for our three algebras follows immediately from the general properties of the theory.

Since we can compute the local cyclic (co)homology of $\mathcal{H}(K)$ directly using Proposition 3.1, we can also use Proposition 5.1 backwards to compute the local cyclic (co)homology of $C^*(K)$ without appealing to the Universal Coefficient Theorem.

6. COMPUTATION OF PERIODIC CYCLIC COHOMOLOGY

The Peter-Weyl Theorem yields that $\mathcal{H}(K)$ is a direct sum of matrix algebras, one for each representation of K . Using Morita invariance and Proposition 3.1, we find that its Hochschild (co)homology vanishes except in dimensions ≥ 1 and

$$\mathrm{HH}_0(\mathcal{H}(K)) \cong R(K) \otimes_{\mathbb{Z}} \mathbb{C}, \quad \mathrm{HH}^0(\mathcal{H}(K)) \cong \mathrm{Hom}_{\mathbb{Z}}(R(K), \mathbb{C}).$$

Thus Connes’s SBI-sequence, which relates Hochschild and cyclic (co)homology, is highly degenerate and yields isomorphisms

$$\begin{aligned} \mathrm{HC}_{2n}(\mathcal{H}(K)) &\cong R(K) \otimes_{\mathbb{Z}} \mathbb{C}, & \mathrm{HC}_{2n+1}(\mathcal{H}(K)) &\cong \{0\}, \\ \mathrm{HC}^{2n}(\mathcal{H}(K)) &\cong \mathrm{Hom}_{\mathbb{Z}}(R(K), \mathbb{C}), & \mathrm{HC}^{2n+1}(\mathcal{H}(K)) &\cong \{0\}, \end{aligned}$$

for all $n \in \mathbb{N}$, the periodicity operator S being the identity. Hence

$$\begin{aligned} \mathrm{HP}_0(\mathcal{H}(K)) &\cong R(K) \otimes_{\mathbb{Z}} \mathbb{C}, & \mathrm{HP}_1(\mathcal{H}(K)) &\cong \{0\}, \\ \mathrm{HP}^0(\mathcal{H}(K)) &\cong \mathrm{Hom}_{\mathbb{Z}}(R(K), \mathbb{C}), & \mathrm{HP}^1(\mathcal{H}(K)) &\cong \{0\}. \end{aligned}$$

As a result,

$$\mathrm{HP}_*(\mathcal{H}(K)) \cong \mathrm{HL}_*(\mathcal{H}(K)) \cong K_*(C^*(K)) \otimes_{\mathbb{Z}} \mathbb{C}$$

and dually for cohomology.

The cyclic type homology theories of $C^\infty(K)$ are similarly easy to compute. This computation is also a trivial special case of results of Victor Nistor and others on the cyclic homology of crossed products. The following lemma asserts that $C^\infty(K)$ is a biprojective Fréchet algebra (see [1, 6]).

Lemma 6.1. *The map*

$$\sigma: C^\infty(K) \rightarrow C^\infty(K) \hat{\otimes} C^\infty(K) \cong C^\infty(K \times K), \quad \sigma f(g_0, g_1) := f(g_0 \cdot g_1),$$

is a $C^\infty(K)$ -bimodule section for the multiplication map

$$m: C^\infty(K) \hat{\otimes} C^\infty(K) \rightarrow C^\infty(K), \quad f_1 \otimes f_2 \mapsto f_1 * f_2.$$

Thus $C^\infty(K)$ is projective as a $C^\infty(K)$ -bimodule.

Its commutator quotient is isomorphic to the space $C^\infty(K/\text{Ad } K)$ of smooth functions on K that are constant on conjugacy classes, that is, invariant with respect to the adjoint action.

Proof. See [9] for the isomorphism $C^\infty(K) \hat{\otimes} C^\infty(K) \cong C^\infty(K \times K)$. It is clear that σ is a well-defined bounded linear map. It is a section for m because the convolution takes the form $m(h)(g) = \int_K h(gg_1, g_1^{-1}) dg_1$ on $C^\infty(K \times K)$. The bimodule structure on $C^\infty(K)$ is defined by left and right convolution, the bimodule structure on $C^\infty(K) \hat{\otimes} C^\infty(K)$ is defined by left convolution on the left and right convolution on the right factor. These actions of $C^\infty(K)$ are integrated forms of group actions of K on these spaces, defined by $\lambda_g f(g_1) = f(g^{-1}g_1)$, $\lambda_g h(g_0, g_1) := h(g^{-1}g_0, g_1)$, $\rho_g f(g_1) = f(g_1g)$ and $\rho_g h(g_0, g_1) := h(g_0, g_1g)$. It is trivial to check that σ is equivariant with respect to these group actions. Hence it is a bimodule map with respect to their integrated forms.

To prove that $C^\infty(K)$ is projective as a bimodule, we iterate σ and consider the corresponding map

$$C^\infty(K) \rightarrow C^\infty(K^3) \subseteq C^\infty(K)^+ \hat{\otimes} C^\infty(K) \hat{\otimes} C^\infty(K)^+.$$

Here $C^\infty(K)^+$ is the space obtained by adjoining a unit to the non-unital algebra $C^\infty(K)$. This map is still a bimodule map and a section for the multiplication map. Since $C^\infty(K)^+ \hat{\otimes} C^\infty(K) \hat{\otimes} C^\infty(K)^+$ is a prototypical free bimodule, $C^\infty(K)$ is projective.

To compute the commutator quotient of $C^\infty(K)$, we consider $C^\infty(K)$ as a direct summand of $C^\infty(K)^+ \hat{\otimes} C^\infty(K)$ via the map $\sigma \circ m$. The commutator quotient of the latter bimodule is canonically isomorphic to $C^\infty(K)$. A computation shows that the map $\sigma \circ m$ induces on commutator quotients the map

$$P: C^\infty(K) \rightarrow C^\infty(K), \quad Pf(g) := \int_K f(h^{-1}gh) dh.$$

This is exactly the projection onto the space of invariants under the adjoint action. Clearly, a smooth function is invariant under the adjoint action if and only if it is constant on conjugacy classes. $\square$

The lemma implies that $C^\infty(K)$ is H -unital (see also [3]). Thus the Hochschild homology of $C^\infty(K)$ can be computed from a projective bimodule resolution of $C^\infty(K)$, we do not have to use $C^\infty(K)^+$. Since $C^\infty(K)$ is itself a projective bimodule, there is a trivial projective resolution. Consequently, Hochschild homology and cohomology vanish in dimensions ≥ 1 and

$$\text{HH}_0(C^\infty(K)) \cong C^\infty(K/\text{Ad } K), \quad \text{HH}^0(C^\infty(K)) \cong C^\infty(K/\text{Ad } K)'.$$

We proceed as for $\mathcal{H}(K)$ to compute the cyclic and periodic cyclic theories and get

$$\begin{aligned} \text{HP}_0(C^\infty(K)) &\cong C^\infty(K/\text{Ad } K), & \text{HP}_1(C^\infty(K)) &\cong \{0\}, \\ \text{HP}^0(C^\infty(K)) &\cong C^\infty(K/\text{Ad } K)', & \text{HP}^1(C^\infty(K)) &\cong \{0\}. \end{aligned}$$

Thus the periodic and local cyclic (co)homology of $C^\infty(K)$ differ and the image of $K_*(C^*(K)) \otimes_{\mathbb{Z}} \mathbb{C}$ under the Chern-Connes character is a proper dense subspace of $\text{HP}_*(C^\infty(K))$.

One can show that the canonical map

$$\mathrm{HP}^0(C^\infty(K)) \rightarrow \mathrm{HL}^0(C^\infty(K)) \cong \prod_{n \in \mathbb{N}} \mathbb{C}$$

is a bijection onto the space of sequences of at most polynomial growth. Therefore, $C^\infty(K/\mathrm{Ad} K) \cong \mathcal{S}$ as a bornological vector space. This follows also from the structure theory of nuclear topological vector spaces because $C^\infty(K/\mathrm{Ad} K)$ is both a subspace and a quotient of $C^\infty(K)$.

To estimate the homological dimension of the algebra $C^\infty(K)$, we have to consider resolutions of $C^\infty(K)^+$, not of $C^\infty(K)$. Nevertheless, biprojectivity implies that the homological dimension is at most 2 (see also [6]).

Lemma 6.2. *The $C^\infty(K)$ -bimodule $C^\infty(K)^+$ has a projective bimodule resolution of length 2. Thus the homological dimension of the algebra $C^\infty(K)$ is at most 2.*

Proof. Since $C^\infty(K)^+$ is an extension of the trivial bimodule $\mathbb{C}$ by the projective bimodule $C^\infty(K)$, it suffices to construct a resolution of length 2 for $\mathbb{C}$. This resolution can then be patched together with the trivial projective resolution of length 0 of $C^\infty(K)$. Abbreviate $A := C^\infty(K)$. The resolution of $\mathbb{C}$ is simply the tensor product of two copies of the extension $A \rightarrow A^+ \rightarrow \mathbb{C}$, that is,

$$\mathbb{C} \leftarrow A^+ \hat{\otimes} A^+ \leftarrow A \hat{\otimes} A^+ \oplus A^+ \hat{\otimes} A \leftarrow A \hat{\otimes} A \leftarrow 0.$$

The bimodule structure on each summand is the obvious one. Since A is projective as a bimodule, it is *a fortiori* projective as a left or right module. Hence this is a projective resolution of the trivial bimodule of length 2. $\square$

We do not really care about the periodic cyclic (co)homology of $C^*(K)$ because we do not expect the periodic theory to yield good results for C^* -algebras, anyway. Nevertheless, we give the result in cohomology because we will use it below and because it is easy to obtain using the amenability of $C^*(K)$. Moreover, since the space of irreducible representations of $C^*(K)$ is 0-dimensional, the result actually is not too bad in this case. Since $C^*(K)$ is a nuclear C^* -algebra, it is an amenable Banach algebra, so that $\mathrm{HH}^n(C^*(K)) = 0$ for $n \geq 1$ (see [7]). The space of bounded traces on $C^*(K)$ is isomorphic to the space $\ell^1(\hat{K}, \mathrm{dim})$, where dim denotes the measure that gives an n -dimensional irreducible representation volume n . Hence

$$\mathrm{HP}^0(C^*(K)) \cong \mathrm{HH}^0(C^*(K)) \cong \ell^1(\hat{K}, \mathrm{dim}), \quad \mathrm{HP}^1(C^*(K)) = \{0\}.$$

We remark that Khalkhali shows that the entire and periodic cyclic cohomology agree for amenable Banach algebras like $C^*(K)$. However, he works with the von Neumann bornology. The story is totally different for the precompact bornology!

7. COMPUTATIONS IN ANALYTIC CYCLIC COHOMOLOGY

In all relevant applications, we have $\mathrm{HA}_* \cong \mathrm{HL}_*$. Such an isomorphism should be expected because $\mathrm{H}_*(\varinjlim C_i) = \varinjlim \mathrm{H}_*(C_i)$, that is, the homology of an inductive system of complexes is already local.

However, the analytic and local theory also involve a completion and the completion in the category of bornological vector spaces may be badly behaved and, in particular, non-local. The problem is that the naïve completion need not be separated, so that we may have to divide out a closure of $\{0\}$. In general, this closure may be surprisingly big and difficult to describe locally. Nevertheless, essentially the only completions we need are completions of tensor products of spaces that are already complete. Usually nothing goes wrong with such completions. Problems can only come from a very serious failure of Grothendieck's approximation property.

If our bornological algebra is nuclear (or, more generally, satisfies the bornological analogue of the approximation property), or if it is a Fréchet algebra equipped with

the precompact bornology, then the completions are automatically separated, so that no problems appear. We will explain this in a forthcoming article.

The algebra $\mathcal{H}(K)$ is nuclear, $C^\infty(K)$ is both nuclear and Fréchet and $C^*(K)$ is a Fréchet algebra. As a result, we have $\mathrm{HA}_* \cong \mathrm{HL}_*$ in these cases, so that

$$\mathrm{HA}_*(\mathcal{H}(K)) \cong \mathrm{HA}_*(C^\infty(K)) \cong \mathrm{HA}_*(C^*(K)) \cong \mathrm{HL}_*(C^*(K)).$$

It is surprising that analytic and periodic cyclic homology are different for $C^\infty(K)$ despite the finite homological dimension.

Since ordinary cohomology is not local, there is no analogue of this in cohomology. In fact, the computation of analytic cyclic cohomology is usually quite difficult. For $\mathcal{H}(K)$, we get through with Proposition 3.1 alone. We see that the periodic, analytic and local theories agree. For the bigger algebras $C^\infty(K)$ and $C^*(K)$, we cannot compute the analytic cyclic cohomology. We only show that it is bigger than the periodic theory. This is straightforward using the Chern-Connes character constructed in [9].

It is well-known that in the entire theory there is a Chern-Connes character for θ -summable Fredholm modules (see [5]). In fact, this was Connes's motivation to introduce the theory. In [9], a different construction is exhibited that produces a Chern-Connes character for arbitrary Fredholm modules without summability conditions, that is, a natural transformation

$$\mathrm{ch}: K^*(A) \rightarrow \mathrm{HA}^*(A)$$

for separable C^* -algebras A . Its composition with the natural map $\mathrm{HA}^*(A) \rightarrow \mathrm{HL}^*(A)$ is the Chern-Connes character in local cyclic cohomology constructed by Puschnigg.

The Chern-Connes character can only exist if A is equipped with the precompact bornology. If A is an amenable Banach algebra, then the result of Khalkhali mentioned above shows that there can be no interesting Chern-Connes character with values in the analytic cyclic cohomology of A equipped with the von Neumann bornology.

Proposition 7.1. *The canonical maps*

$$\mathrm{HA}^*(C^*(K)) \rightarrow \mathrm{HL}^*(C^*(K)) \cong \mathrm{HL}^*(C^\infty(K)) \leftarrow \mathrm{HA}^*(C^\infty(K))$$

are surjective.

Proof. Since $\mathrm{HL}^1 = 0$ in both cases, we only have to consider the even cohomology. The group $K^0(A)$ is isomorphic to the space $\mathrm{Hom}(\mathrm{R}(K), \mathbb{Z})$ by the universal coefficient theorem. We identify this with the group $\mathbb{Z}^{\hat{K}} = \prod_{\rho \in \hat{K}} \mathbb{Z}$ of functions $\hat{K} \rightarrow \mathbb{Z}$. Since analytic cyclic cohomology is a vector space over $\mathbb{C}$, the Chern-Connes character yields a map

$$\mathrm{ch}: \mathbb{Z}^{\hat{K}} \otimes \mathbb{C} \cong K^0(C^*(K)) \otimes \mathbb{C} \rightarrow \mathrm{HA}^0(C^*(K)) \rightarrow \mathrm{HA}^0(C^\infty(K)).$$

If we compose these maps with the natural map to the local cyclic cohomology, we get the canonical map $\mathbb{Z}^{\hat{K}} \otimes \mathbb{C} \rightarrow \prod \mathbb{C} = \mathbb{C}^{\hat{K}}$. It is easy to see that a function $\hat{K} \rightarrow \mathbb{C}$ belongs to its image if and only if its entries are contained in some finitely generated subgroup of $\mathbb{C}$.

Traces on $C^*(K)$ generate another subspace of $\mathrm{HA}^0(C^*(K))$, which is mapped onto the space $\ell^1(\hat{K}, \dim)$. Since $\mathbb{C}$ has dense finitely generated subgroups, any sequence in $\mathbb{C}$ can be decomposed as $(a_n) + (b_n)$ with a sequence (a_n) whose entries lie in a finitely generated subgroup and a sequence $(b_n) \in \ell^1(\hat{K}, \dim)$. Therefore, the map $\mathrm{HA}^0(C^*(K)) \rightarrow \mathrm{HL}^0(C^*(K))$ is surjective. This holds *a fortiori* also for $C^\infty(K)$. $\square$

It is quite plausible that analytic and local cyclic cohomology should agree for $C^*(K)$ or at least for $C^\infty(K)$. However, the author does not know how to prove this. Anyway, the proposition shows that $\mathrm{HA}^*(C^\infty(K))$ is bigger than $\mathrm{HP}^*(C^\infty(K))$, that is, the natural map $\mathrm{HP}^*(C^\infty(K)) \rightarrow \mathrm{HA}^*(C^\infty(K))$ is not surjective. This is striking because $C^\infty(K)$ has finite homological dimension.

8. CONCLUDING REMARKS

If the analytic or local cyclic theories differ from the periodic theory, then they yield results closer to K-theory. This is particularly obvious from the Universal Coefficient Theorem for the local theory. Hence it would be interesting in connection with the Baum-Connes conjecture to compute the local or the analytic cyclic homology for group algebras (both are isomorphic).

It is hard to conceive of a definition of finite dimension for topological or bornological algebras that excludes the biprojective algebra $C^\infty(K)$. Hence we should expect differences between the periodic and analytic theory also for finite dimensional algebras, at least if their K-theory is not finitely generated. It seems plausible that the existence of a finitely summable real spectral triple with sufficiently good properties should guarantee equality between the three cyclic theories. If this spectral triple is a “fundamental class,” we get Poincaré duality between K-theory and K-homology. While this is a reasonable assumption for non-commutative manifolds, it excludes algebras with infinitely generated K-theory like the convolution algebras of compact Lie groups studied above. Hence such a concept of finite dimensionality may not be sufficiently widely applicable.

REFERENCES

- [1] Hanno Baehr, *Stability of continuous cyclic cohomology and operator ideals on Hilbert space*, Ph.D. Thesis, Westfälische Wilhelms-Universität Münster, 2001.
- [2] Bruce Blackadar, *K-theory for operator algebras*, Second, Cambridge University Press, Cambridge, 1998, ISBN 0-521-63532-2. MR **99g**:46104
- [3] Jonathan Block, Ezra Getzler, and John D. S. Jones, *The cyclic homology of crossed product algebras. II. Topological algebras*, J. Reine Angew. Math. **466** (1995), 19–25. MR **96j**:19003
- [4] Alain Connes, *Noncommutative differential geometry*, Inst. Hautes Études Sci. Publ. Math. (1985), 257–360. MR **87i**:58162
- [5] A. Connes, *Entire cyclic cohomology of Banach algebras and characters of θ -summable Fredholm modules*, K-Theory **1** (1988), 519–548. MR **90c**:46094
- [6] A. Ya. Helemskii, *The homology of Banach and topological algebras*, Kluwer Academic Publishers Group, Dordrecht, 1989, ISBN 0-7923-0217-6, Translated from the Russian by Alan West. MR **92d**:46178
- [7] Masoud Khalkhali, *Algebraic connections, universal bimodules and entire cyclic cohomology*, Comm. Math. Phys. **161** (1994), 433–446. MR **95f**:46121
- [8] Jean-Louis Loday, *Cyclic homology*, Second, Springer-Verlag, Berlin, 1998, ISBN 3-540-63074-0. MR **98h**:16014
- [9] Ralf Meyer, *Analytic cyclic cohomology*, Ph.D. Thesis, 1999, arXiv:math.KT/9906205.
- [10] ———, *Excision in entire cyclic cohomology*, J. Eur. Math. Soc. (JEMS) **3** (2001), 269–286. MR **2002g**:46128
- [11] Michael Puschnigg, *Excision in cyclic homology theories*, Invent. Math. **143** (2001), 249–323. MR **2002e**:16014
- [12] ———, *Homotopy functors of ind-algebras and local cyclic cohomology I*, 2002, Preprint, Marseille-Luminy.
- [13] Claude Schochet, *Topological methods for C^* -algebras. III. Axiomatic homology*, Pacific J. Math. **114** (1984), 399–445. MR **86g**:46102

MATHEMATISCHES INSTITUT, WESTFÄLISCHE WILHELMS-UNIVERSITÄT MÜNSTER, EINSTEIN-STR. 62, 48149 MÜNSTER, GERMANY
E-mail address: rameyer@math.uni-muenster.de
URL: <http://www.math.uni-muenster.de/u/rameyer>